

产品名称：盐酸帕罗西汀半水合物

产品别名：Paroxetine hydrochloride hemihydrate

生物活性:	
Description	Paroxetine hydrochloride hemihydrate is a potent selective serotonin-reuptake inhibitor, commonly prescribed as an antidepressant and has GRK2 inhibitory ability with IC ₅₀ of 14 μ M.
IC ₅₀ & Target	IC ₅₀ : 14 μ M (GRK2)[3]
In Vitro	Paroxetine (1 μ M and 10 μ M) distinctly restrains T cell migration induced by CX3CL1 through inhibiting GRK2. Paroxetine inhibits GRK2 induced activation of ERK[1]. Paroxetine (10 μ M) reduces pro-inflammatory cytokines in LPS-stimulated BV2 cells. Paroxetine (0-5 μ M) leads to a dose-dependent inhibition on LPS-induced production of TNF- α and IL-1 β in BV2 cells. Paroxetine also inhibits lipopolysaccharide (LPS)-induced nitric oxide (NO) production and inducible nitric oxide synthase (iNOS) expression in BV2 cells. Paroxetine (5 μ M) blocks LPS-induced JNK activation and attenuates baseline ERK1/2 activity in BV2 cells. Paroxetine relieves microglia-mediated neurotoxicity, and suppresses LPS-stimulated pro-inflammatory cytokines and NO in primary microglial cells[4].
In Vivo	Paroxetine treatment obviously attenuates the symptoms of CIA rats. Paroxetine treatment clearly prevents the histological damage of joints and alleviates T cells infiltration into synovial tissue. Paroxetine reveals a strong effect on inhibiting CX3CL1 production in synovial tissues[1]. Paroxetine (20 mg/kg/day) reduces the myocyte cross-sectional area in rat and ROS formation in the remote myocardium. Paroxetine reduces the susceptibility to ventricular tachycardia. Paroxetine treatment following MI decreases LV remodeling and susceptibility to arrhythmias, probably by reducing ROS formation[2]. In CCI paroxetine-treated group, paroxetine (10 mg/kg, i.p.) produces hyperalgesia at days 7 and 10 ($P < 0.01$), but a decrease in pain behavior is seen at day 14. Moreover, paroxetine (10 mg/kg) significantly attenuates tactile hypersensitivity when compared to CCI vehicle-treated group[5].
References	<p>[1]. Wang Q, et al. Paroxetine alleviates T lymphocyte activation and infiltration to joints of collagen-induced arthritis. <i>Sci Rep</i>. 2017 Mar 28;7:45364.</p> <p>[2]. Liu RP, et al. Paroxetine ameliorates lipopolysaccharide-induced microglia activation via differential regulation of MAPK signaling. <i>J Neuroinflammation</i>. 2014 Mar 12;11:47.</p> <p>[3]. Lassen TR, et al. Effect of paroxetine on left ventricular remodeling in an in vivo rat model of myocardial infarction. <i>Basic Res Cardiol</i>. 2017 May;112(3):26.</p> <p>[4]. Waldschmidt HV, et al. Structure-Based Design of Highly Selective and Potent G Protein-Coupled Receptor Kinase 2 Inhibitors Based on Paroxetine. <i>J Med Chem</i>. 2017 Apr 13;60(7):3052-3069.</p>
实验参考:	
Cell Assay	Cell viability is determined by the tetrazolium salt 3-[4,5-dimethylthiazol-2-yl]-2,5-diphenyltetrazolium bromide (MTT) assay. BV2 and primary microglial cells are initially seeded into 96-well plates at a density of 1×10^4 cells/well and 5×10^4 cells/well, respectively. Following treatment, MTT (5 mg/mL in PBS) is added to each well and incubated at 37°C for four hours. The resulting formazan crystals are dissolved in dimethylsulfoxide (DMSO). The optical density is measured at 570 nm, and results are expressed as a percentage of surviving cells compared with the control. [4]
Animal Administration	Animals are divided into two main groups: 1) pre-emptive and 2) post-injury group. Each main group is divided into three different subgroups: I) CCI vehicle-treated group, II) sham group, and III) CCI paroxetine-treated group. Vehicle is injected i.p. to CCI and sham-operated animals. In the pre-emptive study, paroxetine (10 mg/kg) is injected 1 h before surgery and continued daily until day

	14 post surgery. In the post-injury group, paroxetine (10 mg/kg) is administered at day 7 post injury and continued daily until day 14. All behavioral tests are recorded on day 0 (control day) before surgery and on days 1, 3, 5, 7, 10, and 14 post-nerve injury. [5]
References	<p>[1]. Wang Q, et al. Paroxetine alleviates T lymphocyte activation and infiltration to joints of collagen-induced arthritis. <i>Sci Rep</i>. 2017 Mar 28;7:45364.</p> <p>[2]. Liu RP, et al. Paroxetine ameliorates lipopolysaccharide-induced microglia activation via differential regulation of MAPK signaling. <i>J Neuroinflammation</i>. 2014 Mar 12;11:47.</p> <p>[3]. Lassen TR, et al. Effect of paroxetine on left ventricular remodeling in an in vivo rat model of myocardial infarction. <i>Basic Res Cardiol</i>. 2017 May;112(3):26.</p> <p>[4]. Waldschmidt HV, et al. Structure-Based Design of Highly Selective and Potent G Protein-Coupled Receptor Kinase 2 Inhibitors Based on Paroxetine. <i>J Med Chem</i>. 2017 Apr 13;60(7):3052-3069.</p>



源叶生物