

产品名称: VER-50589

产品别名: VER-50589

生物活性:

Description	VER-50589 is a Hsp90 inhibitor, with an IC ₅₀ of 21 nM and a K _d of 4.5 nM.				
IC ₅₀ & Target	HSP90				
	21 nM (IC ₅₀)				
In Vitro	VER-50589 is a Hsp90 inhibitor, with an IC ₅₀ of 21 nM and a K _d of 4.5 nM. VER-50589 inhibits intrinsic ATPase of full-length recombinant yeast Hsp90, with an IC ₅₀ of 143 ± 23 nM in the presence of 400 μM ATP. VER-50589 shows antiproliferative activities against various human cancer cells, with the lowest GI ₅₀ of 32.7 ± 0.2 nM for CH1 human ovarian cells, and mean GI ₅₀ of 78 ± 15 nM. VER-50589 suppresses the proliferation of human umbilical vein endothelial cells (HUVEC) with GI ₅₀ value of 19 ± 2.4 nM, and shows higher GI ₅₀ s against nontumorigenic human breast (MCF10a) and prostate (PNT2) epithelial cells. Furthermore, VER-50589 displays no differences in cellular activities of isogenic cell lines, and these activities are independent of NQO1 expression. VER-50589 also causes G1 and G2-M block (115 or 575 nM) and induces cytostasis in HCT116 colon cancer cells. In addition, VER-50589 causes great uptake in HCT116 cells[1].				
In Vivo	VER-50589 (4 mg/kg, i.p.) exerts a complete HSP90 inhibition in the athymic mice bearing well-established OVCAR3 human ovarian ascites tumors. VER-50589 (100 mg/kg, i.p.) shows reduced tumor volume and tumor weights in the HCT116 colon carcinoma xenografts compared to the control mice group[1].				
Solvent&Solubility	<i>In Vitro:</i> DMSO : ≥ 48 mg/mL (123.46 mM) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
		1 mM	2.5720 mL	12.8601 mL	25.7202 mL
		5 mM	0.5144 mL	2.5720 mL	5.1440 mL
		10 mM	0.2572 mL	1.2860 mL	2.5720 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液, 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80°C, 6 months; -20°C, 1 month。 -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。				
References	[1]. Sharp SY, et al. Inhibition of the heat shock protein 90 molecular chaperone in vitro and in vivo by novel, synthetic, potent resorcinylc pyrazole/isoxazole amide analogues. Mol Cancer Ther. 2007 Apr;6(4):1198-211.				

实验参考:

Cell Assay	HCT116 and HT29 human colon cancer cells are seeded and left to attach overnight. Vehicle control or compound (VER-50589) is added for 1, 4, and 24 h. Attached cells are collected and counted by hemacytometer. Incubation medium (1 mL) and cell pellets are frozen at -80°C until mass spectrometry analysis[1].
Animal Administration	HCT116 cells (2-5 million cells per site) are injected s.c. in the flanks of 6- to 8-week-old female NCr athymic mice. Dosing commenced when tumors are well established (~5-6 mm diameter). For combined therapy and pharmacokinetic and pharmacodynamic studies, mice bearing HCT116

	xenografts are administered 100 mg/kg VER-50589 i.p. per day for 9 days. Tumor volumes are calculated. On study termination, blood samples are taken, and plasma is separated and stored -80°C. Tumors are excised, weighed, and snap frozen at -80°C[1].
References	[1]. Sharp SY, et al. Inhibition of the heat shock protein 90 molecular chaperone in vitro and in vivo by novel, synthetic, potent resorcinyl pyrazole/isoxazole amide analogues. <i>Mol Cancer Ther.</i> 2007 Apr;6(4):1198-211.



源叶生物