

产品名称: **Cilazapril Monohydrate**
产品别名: 西拉普利一水合物; **Ro 31-2848 monohydrate**

生物活性:				
Description	Cilazapril Monohydrate is a angiotensin-converting enzyme (ACE) inhibitor used for the treatment of hypertension and congestive heart failure. Target: ACE Cilazapril is a new nonthiol group containing angiotensin converting enzyme (ACE) inhibitor. Cilazapril has been investigated in more than 4000 patients with all degrees of hypertension, as well as in the special patient groups such as the elderly, renally impaired, and patients with concomitant diseases, such as congestive cardiac failure or chronic obstructive pulmonary disease [1]. Cilazapril is a very potent and highly effective converting enzyme inhibitor. Doses well below 5 mg/day will probably suffice for therapeutic efficacy [2].			
	In Vitro: DMSO : ≥ 100 mg/mL (229.62 mM) * "≥" means soluble, but saturation unknown.			
Solvent&Solubility	Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg
		1 mM	2.2962 mL	11.4808 mL
		5 mM	0.4592 mL	2.2962 mL
		10 mM	0.2296 mL	1.1481 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液；一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限：-80℃，6 months；-20℃，1 month。-80℃ 储存时，请在 6 个月内使用，-20℃ 储存时，请在 1 个月内使用。 In Vivo: 请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的储备液，再依次添加助溶剂： ——为保证实验结果的可靠性，澄清的储备液可以根据储存条件，适当保存；体内实验的工作液，建议您现用现配，当天使用； 以下溶剂前显示的百分比是指该溶剂在您配制终溶液中的体积占比；如在配制过程中出现沉淀、析出现象，可以通过加热和/或超声的方式助溶 1.请依序添加每种溶剂： 10% DMSO→40% PEG300 →5% Tween-80 → 45% saline Solubility: ≥ 3 mg/mL (6.89 mM); Clear solution 此方案可获得 ≥ 3 mg/mL (6.89 mM，饱和度未知) 的澄清溶液。 以 1 mL 工作液为例，取 100 μL 30.0 mg/mL 的澄清 DMSO 储备液加到 400 μL PEG300 中，混合均匀；向上述体系中加入 50 μL Tween-80，混合均匀；然后继续加入 450 μL 生理盐水定容至 1 mL。 2.请依序添加每种溶剂： 10% DMSO→ 90% (20% SBE-β-CD in saline) Solubility: ≥ 3 mg/mL (6.89 mM); Clear solution 此方案可获得 ≥ 3 mg/mL (6.89 mM，饱和度未知) 的澄清溶液。 以 1 mL 工作液为例，取 100 μL 30.0 mg/mL 的澄清 DMSO 储备液加到 900 μL 20% 的 SBE-β-CD 生理盐水水溶液中，混合均匀。 3.请依序添加每种溶剂： 10% DMSO →90% corn oil Solubility: ≥ 3 mg/mL (6.89 mM); Clear solution 此方案可获得 ≥ 3 mg/mL (6.89 mM，饱和度未知) 的澄清溶液，此方案不适用于实验周期在半个月以上的实			

	验。 以 1 mL 工作液为例，取 100 μL 30.0 mg/mL 的澄清 DMSO 储备液加到 900 μL 玉米油中，混合均匀。
References	[1]. Szucs, T., Cilazapril. A review. Drugs, 1991. 41 Suppl 1: p. 18-24. [2]. Nussberger, J., et al., Repeated administration of the converting enzyme inhibitor cilazapril to normal volunteers. J Cardiovasc Pharmacol, 1987. 9(1): p. 39-44.



源叶生物