



生物活性:					
Description	CC0651 is an allosteric inhibitor of the human Cdc34 ubiquitin-conjugating enzyme. CC0651 potently (IC ₅₀ =1.72 μM) inhibits the ubiquitination of p27 ^{Kip1} , as confirmed by dose-response analysis.				
IC ₅₀ & Target	IC50: 1.72 μM (p27 ^{Kip1} ubiquitination) ^[1]				
In Vitro	CC0651 strongly impairs the rate of ubiquitin chain initiation on substrate by SCF ^{Cdc34} , as measured by the monoubiquitination of Sic1 by K0 ubiquitin. CC0651 actually potentiates the formation of both ubiquitin dimers and monoubiquitinated hCdc34, concordant with the observed accumulation of the hCdc34 conjugate in cells treated with the ester derivative of CC0651. CC0651 completely inhibits the assembly of polyubiquitin chains and decreased formation of free triubiquitin and, to a lesser extent, hCdc34 monoubiquitin, but has no effect on production of diubiquitin ^[1] . CC0651 is an inhibitor of the E2 ubiquitin conjugating enzyme Cdc34A, acts by trapping a weak interaction between ubiquitin and the E2 donor ubiquitin binding site. A quantitative SCF ubiquitination assay with a β-Catenin substrate peptide yields a value of IC ₅₀ of 18±1 μM for CC0651 inhibition, similar to the effective concentrations observed in the NMR and TR-FRET assays ^[2] .				
Solvent&Solubility	In Vitro: DMSO : ≥ 56 mg/mL (126.61 mM) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
		1 mM	2.2610 mL	11.3048 mL	22.6096 mL
		5 mM	0.4522 mL	2.2610 mL	4.5219 mL
		10 mM	0.2261 mL	1.1305 mL	2.2610 mL
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液；一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限：-80℃，6 months；-20℃，1 month。 -80℃ 储存时，请在 6 个月内使用，-20℃ 储存时，请在 1 个月内使用。					
References	[1]. Ceccarelli DF, et al. An allosteric inhibitor of the human Cdc34 ubiquitin-conjugating enzyme. Cell. 2011 Jun 24;145(7):1075-87. [2]. Huang H, et al. E2 enzyme inhibition by stabilization of a low-affinity interface with ubiquitin. Nat Chem Biol. 2014 Feb;10(2):156-63.				
实验参考:					
Cell Assay	PC-3 and HCT116 cell lines are grown in DMEM supplemented with Penicillin/Streptomycin, 2 mM glutamine, and 10% fetal bovine serum (FBS). Lentiviral shRNA constructs for human CDC34, UBE2R2, and nontarget control are used. For proliferation assays, PC-3 cultures are seeded at 5000 cells/well in quadruplicate in 24-well plates, grown for 24 hr, and then infected with lentiviral supernatant. For small-molecule inhibition, cells are seeded in quadruplicate at 500 cells/well in 96-well plates, grown for 24 hr, and treated with compound. Proliferation is measured by MTT				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	assay. PC-3 or HCT 116 cultures are synchronized in G0/G1 by serum starvation for 30 hr and then released by addition of 10% serum. For epistasis experiments, PC-3 cells are infected with CDC34 shRNA for 24 hr, followed by addition of inhibitor and assessment of proliferation after 4 days. Anti-Cdc34, anti-p27, anti-α-tubulin, anti-ubiquitin, and anti-cyclin E antibodies are used^[1].
Kinase Assay	For small-molecule screens, p27^{Kip1} is ubiquitinated in a 15 μL reaction with 40 nM phospho-p27, 40 nM E1, 5 μM E2, 25 nM SCF^{Skp2}, 25 nM Cks1, and 27.8 μM biotinylated ubiquitin in 40 mM Tris-HCl [pH 7.5], 5 mM MgCl₂, 1 mM DTT, and 0.5 mM ATP at 23°C for 3 hr. The reaction is terminated with assay diluent, transferred to 384-well protein A plates coated with 62 ng SC528 antibody, incubated for 1 hr, and washed six times with 10 mM Tris-HCl [pH 7.6], 0.05% Tween20 prior to addition of 25 μL of 0.4 mg/mL europium streptavidin in HEPES [pH 7.6], 1% BSA, 0.2% Tween20 per well. Plates are incubated for 1 hr, washed, and read for Eu-time-resolved fluorescence after addition of 25 μL enhancement solution. For gel-based ubiquitination assays, CC0651 and its analogs are preincubated at the indicated concentrations with 0.5 μg E1, 1–4 μg hCdc34, and 50 ng SCF^{Cdc4}, 100 ng SCF^{Fbw7}, 150 ng SCF^{TrCP}, or 50 ng SCF^{Skp2} for 10–45 min at 4°C in 20 μL reaction buffer (50 mM HEPES [pH 7.5], 10 mM MgCl₂, 2 mM ATP, and 50 μM DTT). Reactions are initiated by addition of 1 μg ubiquitin and respective SCF substrates (50 ng HisSic1 phosphorylated by Cln2-Cdc28, 50 ng cyclin E phosphorylated by Cdk2, 25 ng biotinylated IκBα phosphopeptide [KKERLLDDHDpSGLDpSMKDEE], or 50 ng p27^{Kip1} phosphorylated by cyclin E-Cdk2), incubated at 30°C for 1–3 hr, and products visualized by immunoblot^[1].
References	[1]. Ceccarelli DF, et al. An allosteric inhibitor of the human Cdc34 ubiquitin-conjugating enzyme. Cell. 2011 Jun 24;145(7):1075–87. [2]. Huang H, et al. E2 enzyme inhibition by stabilization of a low-affinity interface with ubiquitin. Nat Chem Biol. 2014 Feb;10(2):156–63.

源叶生物