



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **LEE011 (succinate hydrate)**

产品别名: 瑞博西尼琥珀酸盐水合物; **Ribociclib succinate hydrate**

生物活性:					
Description	Ribociclib succinate hydrate (LEE011 succinate hydrate) is a highly specific CDK4/6 inhibitor with IC ₅₀ values of 10 nM and 39 nM, respectively, and is over 1,000-fold less potent against the cyclin B/CDK1 complex.				
IC ₅₀ & Target	CDK4	CDK6			
	10 nM (IC ₅₀)	39 nM (IC ₅₀)			
In Vitro	Treating a panel of 17 neuroblastoma cell lines with Ribociclib (LEE011) across a four-log dose range (10 to 10,000 nM). Treatment with Ribociclib significantly inhibits substrate adherent growth relative to the control in 12 of the 17 neuroblastoma cell lines examined (mean IC ₅₀ =306±68 nM, considering sensitive lines only, where sensitivity is defined as an IC ₅₀ of less than 1 μM. Ribociclib treatment of two neuroblastoma cell lines (BE2C and IMR5) with demonstrated sensitivity to CDK4/6 inhibition results in a dose-dependent accumulation of cells in the G ₀ /G ₁ phase of the cell cycle. This G ₀ /G ₁ arrest becomes significant at Ribociclib concentrations of 100 nM (p=0.007) and 250 nM (p=0.01), respectively ^[2] .				
In Vivo	CB17 immunodeficient mice bearing BE2C, NB-1643 (MYCN amplified, sensitive in vitro), or EBC1 (non-amplified, resistant in vitro) xenografts are treated once daily for 21 days with Ribociclib (LEE011; 200 mg/kg) or with a vehicle control. This dosing strategy is well tolerated, as no weight loss or other signs of toxicity are observed in any of the xenograft models. Tumor growth is significantly delayed throughout the 21 days of treatment in mice harboring the BE2C or 1643 xenografts (both, p<0.0001), although growth resumed post-treatment ^[2] .				
Solvent&Solubility	In Vitro: DMSO : ≥ 19 mg/mL (33.30 mM) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent Concentration	Mass		
		1 mM	1.7524 mL	8.7621 mL	17.5242 mL
		5 mM	0.3505 mL	1.7524 mL	3.5048 mL
		10 mM	0.1752 mL	0.8762 mL	1.7524 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时, 请在 6 个月内使用, -20℃ 储存时, 请在 1 个月内使用。				
References	[1]. VanArsdale T, et al. Molecular Pathways: Targeting the Cyclin D-CDK4/6 Axis for Cancer Treatment. Clin Cancer Res. 2015 Jul 1;21(13):2905-10. [2]. Rader J, et al. Dual CDK4/CDK6 Inhibition Induces Cell-Cycle Arrest and Senescence in Neuroblastoma. Clin Cancer Res. 2013 Nov 15;19(22):6173-82.				
实验参考:					
		Cells are grown for 24 hours in 35 mm plates, treated with 500 nM Ribociclib for 6 days, and then			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

Cell Assay	fixed and stained overnight. Cells are then imaged for SA- β -gal using an Axio Observer D.1 phase contrast microscope. The percentage of SA- β -gal positive cells is determined by counting the number of positive cells present in three separate microscope frames, and then normalizing to the control. To assess apoptotic activity, cells are plated in triplicate in 96 well plates, treated with Ribociclib, and assayed for caspase 3/7 activation 16 hours after treatment with Caspase-Glo 3/7. Cells treated with SN-38 are used as a positive control ^[2] .
Animal Administration	Mice ^[2] The BE2C, NB-1643, or EBC1 cell line-derived xenografts are implanted subcutaneously into the right flank of CB17 SCID ^{-/-} mice. Animals bearing engrafted tumors of 200-600 mm ³ are then randomized to oral treatment with 200 mg/kg Ribociclib in 0.5 % methylcellulose (n=10) or vehicle (n=10) daily for a total of 21 days. Tumor burden is determined periodically throughout treatment according to the formula $(\pi/6) \times d^2$, where d represents the mean tumor diameter obtained by caliper measurement.
References	[1]. VanArsdale T, et al. Molecular Pathways: Targeting the Cyclin D-CDK4/6 Axis for Cancer Treatment. Clin Cancer Res. 2015 Jul 1;21(13):2905-10. [2]. Rader J, et al. Dual CDK4/CDK6 Inhibition Induces Cell-Cycle Arrest and Senescence in Neuroblastoma. Clin Cancer Res. 2013 Nov 15;19(22):6173-82.

源叶生物