



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **AZD9496 (maleate)**
产品别名: **AZD9496 maleate**

生物活性:

Description	AZD9496 maleate is a potent and selective estrogen receptor (ERα)antagonist with IC ₅₀ of 0.28 nM.				
IC ₅₀ & Target	IC50: 0.28 nM (ERα antagonism), 0.14 nM (ERα downregulation), 0.82 nM (ERα binding) ^[1]				
In Vitro	The potency of AZD9496 with IC ₅₀ of 0.82 nM, 0.14 nM, and 0.28 nM in ERα binding, downregulation, and antagonism, respectively. AZD9496 significantly inhibits MCF-7 cell growth with EC ₅₀ of 0.04 nM ^[1] . Selectivity of AZD9496 over other tested nuclear hormone receptors is high: androgen receptor (AR), IC ₅₀ =30 μM; glucocorticoid receptor (GR), IC ₅₀ =9.2 μM; progesterone receptor (PR), IC ₅₀ =0.54 μM ^[2] .				
In Vivo	Significant tumor growth inhibition is observed as low as 0.5 mg/kg dose in the estrogen-dependent MCF-7 xenograft model, where this effect is accompanied by a dose-dependent decrease in PR protein levels, demonstrating potent antagonist activity. Combining AZD9496 with PI3K pathway and CDK4/6 inhibitors lead to further growth-inhibitory effects compared with monotherapy alone. AZD9496, given once daily orally at 5 and 25 mg/kg produced statistically significant increases in uterine weight compared with the fulvestrant control (P<0.001) but significantly lower than tamoxifen (P=0.001) ^[1] . AZD9496 is also tested in a long-term estrogen deprived model (LTED), using the HCC-1428 LTED cell line that grows in the absence of estrogen and is thought to best represent a model of aromatase inhibition. AZD9496 shows significant activity, with a dose of 5 mg/kg giving tumor regressions in this model ^[2] .				
Solvent&Solubility	In Vitro: DMSO : ≥ 100 mg/mL (179.04 mM) H ₂ O : < 0.1 mg/mL (insoluble) * "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent Concentration Mass	1 mg	5 mg	10 mg
		1 mM	1.7903 mL	8.9517 mL	17.9035 mL
		5 mM	0.3581 mL	1.7904 mL	3.5807 mL
		10 mM	0.1790 mL	0.8952 mL	1.7904 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液, 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时, 请在 6 个月内使用, -20℃ 储存时, 请在 1 个月内使用。 In Vivo: 请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的储备液, 再依次添加助溶剂: ——为保证实验结果的可靠性, 澄清的储备液可以根据储存条件, 适当保存; 体内实验的工作液, 建议您现用现配, 当天使用; 以下溶剂前显示的百分比是指该溶剂在您配制终溶液中的体积占比; 如在配制过程中出现沉淀、析出现象, 可以通过加热和/或超声的方式助溶 1.请依序添加每种溶剂: 10% DMSO→40% PEG300 →5% Tween-80 → 45% saline Solubility: ≥ 2.75 mg/mL (4.92 mM); Clear solution 此方案可获得 ≥ 2.75 mg/mL (4.92 mM, 饱和度未知) 的澄清溶液。				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	以 1 mL 工作液为例, 取 100 μ L 27.5 mg/mL 的澄清 DMSO 储备液加到 400 μ L PEG300 中, 混合均匀向上述体系中加入 50 μ L Tween-80, 混合均匀; 然后继续加入 450 μ L 生理盐水定容至 1 mL。
References	[1]. Weir HM, et al. AZD9496: An Oral Estrogen Receptor Inhibitor That Blocks the Growth of ER-Positive and ESR1-Mutant Breast Tumors in Preclinical Models. Cancer Res. 2016 Jun 1;76(11):3307-18. [2]. De Savi C, et al. Optimization of a Novel Binding Motif to (E)-3-(3,5-Difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenyl)acrylic Acid (AZD9496), a Potent and Orally Bioavailable Selective Estrogen Receptor Downregulator and Antagonist. J Med Chem. 2015 Oct 22;58(20):8128-40.
实验参考:	
Cell Assay	Effect of AZD9496, Fulvestrant, and Tamoxifen on ER α peptide turnover in MCF-7 cells. Cells are grown in steroid-free conditions in SILAC media containing $^{13}\text{C}_6^{15}\text{N}_4$ L-arginine to label ER α peptide as "heavy" (blue line) and then switched to grow in media containing unlabeled L-arginine to label newly synthesized protein as "normal" (red line) with 0.1% DMSO, 300 nM Tamoxifen, 100 nM AZD9496, or 100 nM Fulvestrant for the time indicated. Data shown is representative of two independent experiments ^[1] .
Animal Administration	Mice ^[1] In vivo efficacy of AZD9496 in MCF-7 xenograft model. MCF-7 xenografts, grown in male SCID mice, are dosed daily with either PEG/captisol (vehicle) or AZD9496 (0.02, 0.1, 0.5, 10, and 50 mg/kg, p.o., q.d.). Tumor growth is measured by caliper at regular intervals and mean tumor volumes plotted for each dosed group.
References	[1]. Weir HM, et al. AZD9496: An Oral Estrogen Receptor Inhibitor That Blocks the Growth of ER-Positive and ESR1-Mutant Breast Tumors in Preclinical Models. Cancer Res. 2016 Jun 1;76(11):3307-18. [2]. De Savi C, et al. Optimization of a Novel Binding Motif to (E)-3-(3,5-Difluoro-4-((1R,3R)-2-(2-fluoro-2-methylpropyl)-3-methyl-2,3,4,9-tetrahydro-1H-pyrido[3,4-b]indol-1-yl)phenyl)acrylic Acid (AZD9496), a Potent and Orally Bioavailable Selective Estrogen Receptor Downregulator and Antagonist. J Med Chem. 2015 Oct 22;58(20):8128-40.