



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **AVE 0991 (sodium salt)**
产品别名: **AVE 0991 sodium salt**

生物活性:				
Description	AVE 0991 sodium salt is a nonpeptide and orally active Ang-(1-7) receptor Mas agonist. AVE 0991 competes for high-affinity binding of [¹²⁵ I]-Ang-(1-7) to bovine aortic endothelial cell membranes with IC ₅₀ of 21±35 nM.			
IC ₅₀ & Target	IC50: 21±35 nM (Ang-(1-7) receptor)[1]			
In Vitro	AVE 0991 is a nonpeptide compound that evokes effects similar to Ang-(1-7) on the endothelium. AVE 0991 and unlabeled Ang-(1-7) compete for high-affinity binding of [¹²⁵ I]-Ang-(1-7) to bovine aortic endothelial cell membranes with IC ₅₀ s of 21±35 and 220±280 nM, respectively. Peak concentrations of NO and O ₂ ⁻ release by AVE 0991 sodium salt and Ang-(1-7) (both 10 μM) are not significantly different (NO: 295±20 and 270±25 nM; O ₂ ⁻ : 18±2 and 20±4 nM). However, the released amount of bioactive NO is ≈5 times higher for AVE 0991 in comparison to Ang-(1-7)[1].			
In Vivo	AVE 0991 (0.58 nmol/g) produces a significant decrease of water diuresis in WT mice compared with vehicle-treated animals (0.06±0.03 mL versus 0.27±0.05; n=9 for each group; P<0.01). The antidiuretic effect of AVE 0991 (AVE) is associated with an increase in urine osmolality (1669±231.0 mOsm/KgH ₂ O versus 681.1±165.8 mOsm/KgH ₂ O in vehicle-treated mice; P<0.01). The genetic deletion of <i>Mas</i> abolishes the antidiuretic effect of AVE 0991 during water loading (0.37±0.10 mL [n=9] versus 0.27±0.03 mL [n=11] in AVE 0991-treated mice). As observed with C57BL/6 mice, administration of AVE 0991 (0.58 nmol/g) in water-loaded Swiss mice also produces a significant decrease of the urinary volume compared with vehicle-treated animals (0.13±0.05 mL [n=16] versus 0.51±0.04 mL [n=40]; P<0.01)[2]. One week of treatment with AVE-0991 produces a significant decrease in perfusion pressure (56.55±0.86 vs. 68.73±0.69 mmHg in vehicle-treated rats) and an increase in systolic tension (11.40±0.05 vs. 9.84±0.15 g in vehicle-treated rats), rate of tension rise (+dT/dt; 184.30±0.50 vs. 155.20±1.97 g/s in vehicle-treated rats), rate of tension fall (-dT/dt; 179.60±1.39 vs. 150.80±2.42 g/s in vehicle-treated rats). A slight increase in heart rate (HR) is also observed (220.40±0.71 vs. 214.20±0.74 beats/min in vehicle-treated rats)[3].			
In Vitro: DMSO : ≥ 55 mg/mL (91.26 mM) * "≥" means soluble, but saturation unknown.				
Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
	1 mM	1.6592 mL	8.2960 mL	16.5920 mL
	5 mM	0.3318 mL	1.6592 mL	3.3184 mL
	10 mM	0.1659 mL	0.8296 mL	1.6592 mL
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液; 一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80°C, 6 months; -20°C, 1 month。 -80°C 储存时, 请在 6 个月内使用, -20°C 储存时, 请在 1 个月内使用。 In Vivo: 请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的使				



Solvent&Solubility	备液, 再依次添加助溶剂: ——为保证实验结果的可靠性, 澄清的储备液可以根据储存条件, 适当保存; 体内实验的工作液, 建议您现用现配, 当天使用; 以下溶剂前显示的百分比是指该溶剂在您配制终溶液中的体积占比; 如在配制过程中出现沉淀、析出现象, 可以通过加热和/或超声的方式助溶 1.请依序添加每种溶剂: 10% DMSO→40% PEG300 →5% Tween-80 → 45% saline Solubility: ≥ 2.08 mg/mL (3.45 mM); Clear solution 此方案可获得 ≥ 2.08 mg/mL (3.45 mM, 饱和度未知) 的澄清溶液。 以 1 mL 工作液为例, 取 100 μL 20.8 mg/mL 的澄清 DMSO 储备液加到 400 μL PEG300 中, 混合均匀向上述体系中加入 50 μL Tween-80, 混合均匀; 然后继续加入 450 μL 生理盐水定容至 1 mL。 2.请依序添加每种溶剂: 10% DMSO→ 90% (20% SBE-β-CD in saline) Solubility: ≥ 2.08 mg/mL (3.45 mM); Clear solution 此方案可获得 ≥ 2.08 mg/mL (3.45 mM, 饱和度未知) 的澄清溶液。 以 1 mL 工作液为例, 取 100 μL 20.8 mg/mL 的澄清 DMSO 储备液加到 900 μL 20% 的 SBE-β-CD 生理盐水水溶液中, 混合均匀。 3.请依序添加每种溶剂: 10% DMSO →90% corn oil Solubility: ≥ 2.08 mg/mL (3.45 mM); Clear solution 此方案可获得 ≥ 2.08 mg/mL (3.45 mM, 饱和度未知) 的澄清溶液, 此方案不适用于实验周期在半个月以上的实验。 以 1 mL 工作液为例, 取 100 μL 20.8 mg/mL 的澄清 DMSO 储备液加到 900 μL 玉米油中, 混合均匀。
References	[1]. Wiemer G, et al. AVE 0991, a nonpeptide mimic of the effects of angiotensin-(1-7) on the endothelium. Hypertension. 2002 Dec;40(6):847-52. [2]. Pinheiro SV, et al. Nonpeptide AVE 0991 is an angiotensin-(1-7) receptor Mas agonist in the mouse kidney. Hypertension. 2004 Oct;44(4):490-6. [3]. Ferreira AJ, et al. The nonpeptide angiotensin-(1-7) receptor Mas agonist AVE-0991 attenuates heart failure induced by myocardial infarction. Am J Physiol Heart Circ Physiol. 2007 Feb;292(2):H1113-9.
实验参考:	
Animal Administration	Mice[2] Swiss male mice, Mas-KO (Mas^{-/-}) male mice on the pure genetic background C57BL/6, and WT C57BL/6 control mice (Mas^{+/+}) are used. Water diuresis is induced by intraperitoneal water injection (0.05 mL/g of body weight [BW]) in conscious mice. Drugs are administered in the same injection with water load at prefixed volumes (0.01 mL/g BW). In the first set of experiments, WT mice (C57BL/6, control group) or Mas-KO mice are treated with: (1) 0.58 nmol/g AVE 0991 (n=9, control; n=11, Mas-KO mice); or (2) vehicle for AVE 0991 (10 μM KOH, 0.01 mL/g; n=9, control; n=9, Mas-KO). In the second set, Swiss mice are treated with: (1) vehicle (n=36); (2) 0.58 nmol/g AVE 0991 (n=16); (3) 46 pmol/g Ang-(1-7) antagonist A-779 (n=4); (4) 2 nmol/g losartan or valsartan (n=5); (5) 2 nmol/g AT₂ receptor antagonists PD123319 or PD123177 (n=9); (6) AVE 0991 combined with A-779; (7) AVE 0991 combined with losartan or valsartan (n=4 for each); (8) or AVE 0991 combined with PD123319 (n=5) or PD123177 (n=4). The urinary volume is measured for 60



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	minutes after water loading, and urine samples are obtained to determine the osmolality. The dose of AVE 0991 is based in preliminary experiments performed in Swiss mice. Rats[3] Male Wistar rats weighting 250-300 g are used. Rats are treated either with AVE-0991 (1 mg/kg, n=9) or vehicle (0.9% NaCl, n=11) administered orally by gavage. At the end of the 7 day period of AVE-0991 treatment, the animals are decapitated 10-15 min after intraperitoneal injection of 400 IU of heparin. After the thorax is opened, the heart is carefully dissected, removed from the thoracic cavity, and placed in a plate containing ice-cold Krebs-Ringer solution (KRS) to attenuate any potential cardiac damage during dissection of aorta artery.
Kinase Assay	Binding of [¹²⁵I]-Ang-(1-7) is performed. Briefly, 100 µg of membranes from primary cultured bovine aortic endothelial cells (BAECs, passage 1) are incubated in a total volume of 200 µL for 45 minutes at 25°C in HEPES-buffered saline (10 mM HEPES, 0.1 M NaCl, 5 mM MgCl₂) containing 0.2% BSA and protease inhibitor cocktail Complete (Boehringer Mannheim). Saturable binding of [¹²⁵I]-Ang-(1-7) is calculated by subtracting nonspecific binding (40% to 50%), determined in the presence of 10 µM unlabeled Ang-(1-7) from total binding. Competition experiments with increasing concentrations of AVE 0991 and unlabeled Ang-(1-7) are performed in the presence of 10 nM [¹²⁵I]-Ang-(1-7). Assays are terminated by vacuum filtration (≤15 mm Hg) over Durapore filters (0.65 µm, Opak 96-well plates) presoaked with 1% BSA. The filters are washed 3 times with each 100 µL of PBS (50 mM, NaHPO₄ and 0.15 M NaCl, pH 7.2). Radioactivity on dried filters is quantified with a gamma counter[1].
References	[1]. Wiemer G, et al. AVE 0991, a nonpeptide mimic of the effects of angiotensin-(1-7) on the endothelium. Hypertension. 2002 Dec;40(6):847-52. [2]. Pinheiro SV, et al. Nonpeptide AVE 0991 is an angiotensin-(1-7) receptor Mas agonist in the mouse kidney. Hypertension. 2004 Oct;44(4):490-6. [3]. Ferreira AJ, et al. The nonpeptide angiotensin-(1-7) receptor Mas agonist AVE-0991 attenuates heart failure induced by myocardial infarction. Am J Physiol Heart Circ Physiol. 2007 Feb;292(2):H1113-9.