



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **Vapreotide (acetate)**

产品别名: **醋酸伐普肽 ; RC-160 acetate; BMY-41606 acetate**

生物活性:				
Description	Vapreotide acetate (RC-160 acetate; BMY-41606 acetate) is a neurokinin-1 (NK1) receptor antagonist, with an IC50 of 330 nM.			
In Vitro	Vapreotide attenuates the Substance P (SP)-triggered intracellular calcium increases and NF-κB activation in a dose-dependent manner. Vapreotide also inhibits SP-induced IL-8 and MCP-1 production in HEK293-NK1R and U373MG cell lines. Vapreotide inhibits HIV-1 infection of human MDM <i>in vitro</i> , an effect that is reversible by SP pretreatment[1]. Vapreotide significantly inhibits GH-, PRL, and/or alpha-subunit release by human GH-secreting pituitary adenoma cells in concentrations as low as 10 ⁻¹² -10 ⁻¹⁴ M. Vapreotide inhibits GH release with an IC ₅₀ of 0.1 pM[2]. Vapreotide exhibits moderate-to-high affinities for SSTR2, -3, and -5 (IC ₅₀ =0.17, 0.1 and 21 nM, respectively) and low affinity for SSTR1 and -4 (IC ₅₀ =200 and 620 nM, respectively). RC-160 inhibits serum-induced proliferation of CHO cells expressing SSTR2 and SSTR5 (EC ₅₀ =53 and 150 pM, respectively)[3].			
In Vivo	In cirrhosis, bleeding by rupture of oesophagogastric varices is a severe complication of portal hypertension. The acute administration of vapreotide to rats decreases collateral circulation blood flow while chronic administration attenuated its development[4]. Tumor volumes and weights are reduced by about 40% by RC-160 by s.c. injections at doses of 100 µg/day/animal. Vapreotide can inhibit the growth of androgen-independent prostate cancer when the therapy is started at an early stage of tumor development[5].			
Solvent&Solubility	In Vitro: H ₂ O : 14.29 mg/mL (11.99 mM; Need ultrasonic)			
		Solvent Mass Concentration	1 mg	5 mg
	Preparing	1 mM	0.8394 mL	4.1968 mL
	Stock Solutions	5 mM	0.1679 mL	0.8394 mL
		10 mM	0.0839 mL	0.4197 mL
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时，请在 6 个月内使用， -20℃ 储存时，请在 1 个月内使用。				
References	[1]. Spitsin S, et al. Analog of somatostatin vapreotide exhibits biological effects in vitro via interaction with neurokinin-1 receptor. Neuroimmunomodulation. 2013;20(5):247-55. [2]. Hofland LJ, et al. Relative potencies of the somatostatin analogs octreotide, BIM-23014, and RC-160 on the inhibition of hormone release by cultured human endocrine tumor cells and normal rat anterior pituitary cells. Endocrinology. 1994 Jan;134(1):301-6. [3]. Buscail L, et al. Inhibition of cell proliferation by the somatostatin analogue RC-160 is mediated by somatostatin receptor subtypes SSTR2 and SSTR5 through different mechanisms. Proc Natl Acad Sci U S A. 1995 Feb 28;92(5):1580-4.			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	[4]. Veal N, et al. Hemodynamic effects of acute and chronic administration of vapreotide in rats with cirrhosis. Dig Dis Sci. 2003 Jan;48(1):154-61. [5]. Pinski J, et al. Effect of somatostatin analog RC-160 and bombesin/gastrin releasing peptide antagonist RC-3095 on growth of PC-3 human prostate-cancer xenografts in nude mice.
实验参考:	
Cell Assay	CHO cells are cultured in aMEM containing 10% FCS. After overnight attachment, the medium is changed to a MEM containing either 10% FCS or insulin with and without vapreotide. Cell growth is measured after 24 h by counting cells with a Coulter counter model ZM[3].
Animal Administration	Rats: Acute effects are evaluated at baseline and 30 min after placebo or vapreotide (8 µg/kg/hr) infusions in DMNA rats. Chronic hemodynamic effects are evaluated using subcutaneous implants for five weeks in anesthetized DMNA rats and in sham rats. Hemodynamic measurements include splenorenal shunt blood flow by the transit time ultrasound method and cardiac output by the combined dilution-TTU method[4]. Mice: Nude mice bearing xenografts of the androgen-independent human prostate-cancer cell line PC-3 are treated for 4 weeks with somatostatin analog vapreotide (20 µg/day/animal), bombesin/gastrin-releasing peptide (GRP) antagonist, or the combination of both peptides. Tumor volumes and weights are measured[5].
References	[1]. Spitsin S, et al. Analog of somatostatin vapreotide exhibits biological effects in vitro via interaction with neurokinin-1 receptor. Neuroimmunomodulation. 2013;20(5):247-55. [2]. Hofland LJ, et al. Relative potencies of the somatostatin analogs octreotide, BIM-23014, and RC-160 on the inhibition of hormone release by cultured human endocrine tumor cells and normal rat anterior pituitary cells. Endocrinology. 1994 Jan;134(1):301-6. [3]. Buscail L, et al. Inhibition of cell proliferation by the somatostatin analogue RC-160 is mediated by somatostatin receptor subtypes SSTR2 and SSTR5 through different mechanisms. Proc Natl Acad Sci U S A. 1995 Feb 28;92(5):1580-4. [4]. Veal N, et al. Hemodynamic effects of acute and chronic administration of vapreotide in rats with cirrhosis. Dig Dis Sci. 2003 Jan;48(1):154-61. [5]. Pinski J, et al. Effect of somatostatin analog RC-160 and bombesin/gastrin releasing peptide antagonist RC-3095 on growth of PC-3 human prostate-cancer xenografts in nude mice.