



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **NBI-74330**
产品别名: **NBI-74330**

生物活性:					
Description		NBI-74330 is a potent antagonist for CXCR3 , and exhibits potent inhibition of (¹²⁵ I)CXCL10 and (¹²⁵ I)CXCL11 specific binding with K_i of 1.5 and 3.2 nM, respectively.			
IC ₅₀ & Target		[¹²⁵ I]CXCL11-CXCR3			
		[¹²⁵ I]CXCL10-CXCR3			
		1.5 nM (K _i in CXCR3-CHO cell membranes)			
		3.2 nM (K _i in CXCR3-CHO cell membranes)			
In Vitro		BI-74330 demonstrates potent inhibition of [¹²⁵ I]CXCL11 specific binding to membranes prepared from transfected CHO cells expressing CXCR3 (CXCR3-CHO) (K _i =3.6 nM). NBI-74330 is 12- and 3.5-fold more potent than CXCL9 (K _i =45.2 nM) and CXCL10 (K _i =12.5 nM), respectively, at inhibiting [¹²⁵ I]CXCL11 binding to CXCR3-CHO cell membranes. NBI-74330 inhibits calcium mobilization in response to CXCL11 and CXCL10 with an IC ₅₀ value of 7 nM for both ligands used at their EC ₈₀ concentrations (1 nM for CXCL11 and 30 nM for CXCL10). NBI-74330 specifically inhibits CXCR3-mediated calcium mobilization. NBI-74330 also dose-dependently inhibits CXCL11-induced [³⁵ S]GTPγS binding in membranes of cells endogenously expressing CXCR3 (H9 cells, IC ₅₀ value 5.5 nM). BI-74330 inhibits CXCL11-induced chemotaxis in these cells with an IC ₅₀ of 3.9 nM[1]. NBI-74330 (30-300 nm, 1-10 μM) produces concentration-dependent, parallel rightward shifts of the CXCL11 E/[A] curve with no significant change in the E/[A] curve maximal response[2].			
In Vivo		NBI-74330 (100 mg/kg) results in the formation of an N-oxide metabolite, also an antagonist of CXCR3, in mice[2]. Mice treated with 100 mg/kg NBI-74330 (in 1% Na Doc in 0.5% 400Cp Methylcellulose) result in serum concentrations of approximately 1 μM. This concentration is sufficient to fully block the CXCR3 receptor in vivo[3].			
Solvent&Solubility		In Vitro: DMSO : ≥ 35 mg/mL (57.80 mM) * "≥" means soluble, but saturation unknown.			
		Solvent Mass Concentration 1 mg 5 mg 10 mg			
		Preparing			
		Stock Solutions			
		*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限：-80℃，6 months；-20℃，1 month。 -80℃ 储存时，请在 6 个月内使用，-20℃ 储存时，请在 1 个月内使用。			
		[1]. Heise CE, et al. Pharmacological characterization of CXC chemokine receptor 3 ligands and a small molecule antagonist. J Pharmacol Exp Ther. 2005 Jun;313(3):1263-71. [2]. Jopling LA, et al. Analysis of the pharmacokinetic/pharmacodynamic relationship of a small molecule			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

References	CXCR3 antagonist, NBI-74330, using a murine CXCR3 internalization assay. Br J Pharmacol. 2007 Dec;152(8):1260-71. [3]. van Wanrooij EJ, et al. CXCR3 antagonist NBI-74330 attenuates atherosclerotic plaque formation in LDL receptor-deficient mice. Arterioscler Thromb Vasc Biol. 2008 Feb;28(2):251-7.
实验参考:	
Animal Administration	Female LDLR^{-/-} mice, 10 weeks old (n=8 to 12 per group), are fed a Western-type diet containing 0.25% cholesterol and 15% cocoa butter 2 weeks before collar placement. 20 Mice are treated with a subcutaneous injection of 100 mg/kg NBI-74330 every day during the entire experiment. After 8 weeks of Western-type diet and treatment, the mice are euthanized and organs are harvested for histology, fluorescence-activated-cell sorter (FACS) analysis, and RNA isolation. Blood samples are collected by tail bleeding from nonfasted animals, and concentrations of serum cholesterol and triglycerides are determined using enzymatic colorimetric procedures. [3]
Kinase Assay	Cell membrane fractions are resuspended in 50 mM HEPES, 10 mM MgCl₂, 100 mM NaCl, and 1 mM CaCl₂, pH 7.2 for use in competitive radioligand binding reactions. Reactions are performed in duplicate and consisted of 25-μL unlabeled chemokine at indicated concentrations, 25-μL radiolabeled chemokine ligand (appr 70 nM; [¹²⁵I]CXCL11 and [¹²⁵I]CXCL10 with specific activities of 1500 and 2200 Ci/mmol, respectively; Fifty-μL membrane protein (5 μg) are added sequentially in assay buffer (50 mM HEPES, 10 mM MgCl₂, 100 mM NaCl, 1 mM CaCl₂, and 0.1% BSA, pH 7.2) to low-binding 96-well plates. The reaction is allowed to reach equilibrium by incubation at room temperature for 45 min while shaking. The amount of bound radioligand is determined by harvesting membranes via filtration through a UniFilter GF/C filter plate using a UniFilter-96 vacuum manifold (filters are pretreated with 1% polyethylenimine), washing twice with 400-μL wash buffer (10 mM HEPES, 5 mM MgCl₂, 1 mM CaCl₂, and 500 mM NaCl, pH 7.3), and measuring radioactivity by liquid scintillation using a TopCount NXT. The dissociation half-life of [¹²⁵I]CXCL11 is measured using CXCR3-CHO membranes that are allowed to equilibrate with radiolabel (appr 70 nM) for 30 min prior to the addition of excess cold CXCL11 (31 nM final) in the presence or absence of different concentrations of NBI-74330. Membrane-bound [¹²⁵I]CXCL11 is assessed in duplicate along with nonspecific binding ([¹²⁵I]CXCL11 plus excess cold CXCL11 added at the same) and total binding ([¹²⁵I]CXCL11 without inhibitors) at each time point on the same plate. [1]
References	[1]. Heise CE, et al. Pharmacological characterization of CXCR3 chemokine receptor 3 ligands and a small molecule antagonist. J Pharmacol Exp Ther. 2005 Jun;313(3):1263-71. [2]. Jopling LA, et al. Analysis of the pharmacokinetic/pharmacodynamic relationship of a small molecule CXCR3 antagonist, NBI-74330, using a murine CXCR3 internalization assay. Br J Pharmacol. 2007 Dec;152(8):1260-71. [3]. van Wanrooij EJ, et al. CXCR3 antagonist NBI-74330 attenuates atherosclerotic plaque formation in LDL receptor-deficient mice. Arterioscler Thromb Vasc Biol. 2008 Feb;28(2):251-7.