



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **EMD534085**
产品别名: **EMD534085**

生物活性:

Description	EMD534085 is a potent and selective inhibitor of the mitotic kinesin-5 with an IC50 of 8 nM.				
IC50 & Target	Kinesin-5				
	8 nM (IC50)				
In Vitro	EMD 534085 does not inhibit any other tested kinesins (BimC, CEN-PE, Chromokinesin, KHC, KIF3C, KIFC3, MKLP-1, and MCAK) at 1 μM or 10 μM concentration, showing selectively over kinesin-5. EMD 534085 binds to the allosteric pocket of kinesin-5[1]. EMD534085 induces rapid cell death in HL60 during mitotic arrest. Caspase-8, -9, -3, -7 are activated; Parp1 is cleaved; Mcl1 and XIAP are degraded in EMD534085-treated HL60 cells. EMD534085 treated HL60 cells also shows significantly accumulated phospho-Histone H3 level starting at 6 hrs post thymidine release[2].				
In Vivo	In a low dose PK of EMD 534085 in mice the clearance is 1.8 L/h/kg on average, the volume of distribution is 7.4 L/kg and the half life around 2.5 h. The bioavailability in high dose experiments (>10 mg/kg) is always above 50% in mice. Intraperitoneal administration of EMD 534085 enables significant systemic exposure in mice leading to a significant tumor growth reduction without toxic side effects[1].				
Solvent&Solubility	<i>In Vitro:</i>				
	DMSO : ≥ 26 mg/mL (54.56 mM)				
	* "≥" means soluble, but saturation unknown.				
	Preparing Stock Solutions	Solvent / Mass / Concentration	1 mg	5 mg	10 mg
		1 mM	2.0985 mL	10.4925 mL	20.9850 mL
		5 mM	0.4197 mL	2.0985 mL	4.1970 mL
10 mM		0.2099 mL	1.0493 mL	2.0985 mL	
*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限 -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时，请在 6 个月内使用， -20℃ 储存时，请在 1 个月内使用。					
References	[1]. Schiemann K, et al. The discovery and optimization of hexahydro-2H-pyrano[3,2-c]quinolines (HHPQs) as potent and selective inhibitors of the mitotic kinesin-5. Bioorg Med Chem Lett. 2010 Mar 1;20(5):1491-5. [2]. Tang Y, et al. Rapid induction of apoptosis during Kinesin-5 inhibitor-induced mitotic arrest in HL60 cells. Cancer Lett. 2011 Nov 1;310(1):15-24.				
实验参考:					
Cell Assay	Epithelial cell lines HeLa and MCF7 are synchronized in G2-phase using RO-3306. Cells are treated with 10 μM RO-3306 for 16 hrs, and then are washed and released to either warm growth medium or medium supplemented with 500 nM EMD534085[2].				
	[1]. Schiemann K, et al. The discovery and optimization of hexahydro-2H-pyrano[3,2-c]quinolines (HHPQs)				



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

References	as potent and selective inhibitors of the mitotic kinesin-5. Bioorg Med Chem Lett. 2010 Mar 1;20(5):1491-5. [2]. Tang Y, et al. Rapid induction of apoptosis during Kinesin-5 inhibitor-induced mitotic arrest in HL60 cells. Cancer Lett. 2011 Nov 1;310(1):15-24.
-------------------	--



源叶生物