



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

产品名称: **Exendin-4 (Acetate)**
产品别名: **醋酸艾塞那肽; Exenatide acetate**

生物活性:				
Description	Exendin-4 Acetate (Exenatide acetate), a 39 amino acid peptide, is a long-acting glucagon-like peptide-1 receptor agonist with an IC ₅₀ of 3.22 nM.			
IC₅₀ & Target	IC ₅₀ : 3.22 nM (glucagon-like peptide-1 receptor)[1]			
In Vitro	In human umbilical vein endothelial cells, Exendin-4 significantly increases NO production, endothelial NO synthase (eNOS) phosphorylation, and GTP cyclohydrolase 1 (GTPCH1) level in a dose-dependent manner[2]. Exendin-4 shows cytotoxic effects to MCF-7 breast cancer cells with IC ₅₀ of 5 µM at 48 hour[3].			
In Vivo	Both low- and high-dose Exendin-4 treatment in ob/ob mice improve serum ALT and reduce serum glucose, and calculated HOMA scores compared with control. Exendin-4-treated ob/ob mice sustain a marked reduction in the net weight gain in the final 4 weeks of the study period[4]. Animals treated with Exendin-4 have more pancreatic acinar inflammation, more pyknotic nuclei and weigh significantly less than control rats. Exendin-4 treatment is associated with lower leptin levels as well as lower HOMA values in rats[5]. Exenatide causes dose-dependent relaxation of rat thoracic aorta, which is evoked via the GLP-1 receptor and is mediated mainly by H ₂ S but also by NO and CO[6].			
Solvent&Solubility	<i>In Vitro:</i> DMSO : ≥ 66.66 mg/mL (15.70 mM) H₂O : 1 mg/mL (0.24 mM; Need ultrasonic) * "≥" means soluble, but saturation unknown.			
	Preparing Stock Solutions	Solvent Mass Concentration	1 mg	5 mg
		1 mM	0.2355 mL	1.1774 mL
		5 mM	0.0471 mL	0.2355 mL
		10 mM	0.0235 mL	0.1177 mL
	*请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液。一旦配成溶液, 请分装保存, 避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80℃, 6 months; -20℃, 1 month。 -80℃ 储存时, 请在 6 个月内使用, -20℃ 储存时, 请在 1 个月内使用。			
References	[1]. Doyle ME, et al. The importance of the nine-amino acid C-terminal sequence of exendin-4 for binding to the GLP-1 receptor and for biological activity. Regul Pept. 2003 Jul 15;114(2-3):153-8. [2]. Wei R, et al. Exenatide exerts direct protective effects on endothelial cells through the AMPK/Akt/eNOS pathway in a GLP-1 receptor-dependent manner. Am J Physiol Endocrinol Metab. 2016 Jun 1;310(11):E947-57. [3]. Fidan-Yaylall G, et al. Antidiabetic exendin-4 activates apoptotic pathway and inhibits growth of breast cancer cells. Tumour Biol. 2016 Feb;37(2):2647-53. [4]. Ding X, et al. Exendin-4, a glucagon-like protein-1 (GLP-1) receptor agonist, reverses hepatic steatosis in ob/obmice. Hepatology. 2006 Jan;43(1):173-81.			



上海源叶生物科技有限公司
Shanghai yuanye Bio-Technology Co., Ltd
电话: 021-61312973 传真: 021-55068248
网址: www.shyuanye.com
邮箱: shyysw@sina.com

	[5]. Nachnani JS, et al. Biochemical and histological effects of exendin-4 (exenatide) on the rat pancreas. Diabetologia. 2010 Jan;53(1):153-9. [6]. Sélley E, et al. Exenatide induces aortic vasodilation increasing hydrogen sulphide, carbon monoxide and nitric oxide production. Cardiovasc Diabetol. 2014 Apr 2;13:69.
实验参考:	
Animal Administration	Rats: 20 Sprague-Dawley male rats, ten of which are treated with exendin-4 (10 µg/kg) and ten of which are used as controls. The study period is 75 days. Serum and pancreatic tissue are removed for biochemical and histological study. Blood glucose, amylase, lipase and adipocytokines are compared between the two groups[5]. Mice: The exendin-4 treatment groups are treated with 10 µg/kg every 24 hours for the first 14 days. This treatment is the induction phase. Respective control mice (lean and ob/ob) receive saline every 24 hours. After 14 days Exendin-4-treated mice are randomly divided into two groups: one group receives high dose exendin-4 (20 µg/kg) every 12 hours, while the second group continues with low dose exendin-4 (10 µg/kg) every 12 hours. The control mice continue to receive saline every 12 hours. The mice are weighed daily for the 60-day treatment period[4].
References	[1]. Doyle ME, et al. The importance of the nine-amino acid C-terminal sequence of exendin-4 for binding to the GLP-1 receptor and for biological activity. Regul Pept. 2003 Jul 15;114(2-3):153-8. [2]. Wei R, et al. Exenatide exerts direct protective effects on endothelial cells through the AMPK/Akt/eNOS pathway in a GLP-1 receptor-dependent manner. Am J Physiol Endocrinol Metab. 2016 Jun 1;310(11):E947-57. [3]. Fidan-Yaylali G, et al. Antidiabetic exendin-4 activates apoptotic pathway and inhibits growth of breast cancer cells. Tumour Biol. 2016 Feb;37(2):2647-53. [4]. Ding X, et al. Exendin-4, a glucagon-like protein-1 (GLP-1) receptor agonist, reverses hepatic steatosis in ob/obmice. Hepatology. 2006 Jan;43(1):173-81. [5]. Nachnani JS, et al. Biochemical and histological effects of exendin-4 (exenatide) on the rat pancreas. Diabetologia. 2010 Jan;53(1):153-9. [6]. Sélley E, et al. Exenatide induces aortic vasodilation increasing hydrogen sulphide, carbon monoxide and nitric oxide production. Cardiovasc Diabetol. 2014 Apr 2;13:69.